Appendix A: PrEP Algorithm (Front of Card)

Men who have Sex with Men/Transgender Women

Condomless anal sex in last 6 months with:

- Infectious syphilis or rectal bacterial STI in last year
- Recurrent use of post-exposure prophylaxis (PEP)
- Ongoing sexual relationship with HIV+ partner with substantial transmission risk
- HRI-MSM score ≥ 11 (compelling indication)

Not indicated: Serodiscordant, closed relationships where HIV+ partner is on ART and negligible risk of HIV transmission (suppressed viral load and no bacterial STIs)

Heterosexuals

Condomless anal/vaginal sex in serodiscordant relationship where HIV+ partner has significant (also consider if non-negligible) transmission risk

Consider in high-risk pregnant women

Adults with multiple partners of unknown status – decisionbased on local HIV risk and patient risk behaviours

Not indicated: Serodiscordant, closed relationships where HIV+ partner is on ART and negligible risk of HIV transmission (suppressed viral load and no bacterial STIs)

People Who Inject Drugs


If there is risk of HIV infection

 Assess risk based on local epidemiology

Note: Ensure other harm reduction strategies have been implemented if available (clean injection equipment, opioid substitution therapy, etc.)

Initial Assessment (See Baseline Lab Evaluations)

- · Negative HIV test in last week
- Evaluate for signs or symptoms suggestive of acute HIV infection in last 12 weeks
- If acute HIV suspected, delay PrEP and repeat test in 7-21 days
- eGFR ≥ 60mL/min
- Counsel patient on other methods of risk reduction and adverse effects
- · Determine affordability



SK 🥵 HIV
COLLABORATIVE

PrEP Algorithm

January 2018